SOLAR OBSERVATIONS

SOLAR RADIATION MEASUREMENTS DURING MARCH 1933

By IRVING F. HAND, Assistant in Solar Radiation Investigations

For a description of instruments and their exposures, the reader is referred to the January 1932 Review, page 26.

Table 1 shows that solar radiation intensities averaged above normal for February at Washington and Madison,

and close to normal at Lincoln.

Table 2 shows an excess in the total solar radiation received on a horizontal surface at all stations except Madison, Lincoln, Pittsburgh, and La Jolla.

Table 3 shows the effect of cirri on turbidity factors;

particularly on the 27th.

Polarization measurements made on 5 days at Washington give a mean of 49 percent with a maximum of 54 percent on the 29th. These are slightly below normal for the month. No polarization readings were obtained at Madison due to the presence of ice and snow.

Table 1.—Solar radiation intensities during March 1933 [Gram-calories per minute per square centimeter of normal surface] Washington, D.C.

	Sun's zenith distance.												
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon		
me	75th	Air mass											
	mer. time		A.N	1.			P.M.				solar time		
	е.	5.0	4.0	3.0	2.0	1 1.0	2.0	3.0	4.0	5.0	e		
Mar. 6 Mar. 8 Mar. 9	mm 2. 62 5. 16 2. 11	cal. 0.87 .83			cal. 1. 29 1. 32	cal.	cal.	cal.	cal.	cal.	mm 1. 45 4. 57 1. 24		
Mar. 17 Mar. 27 Mar. 28	3. 81 3. 15 4. 17		. 59	. 77	. 88						4. 75 2. 74 3. 99		
Mar. 29 Means Departures	2. 74	.80 .74 +.01	.91	1.03 1.04 +0.09	1.19	1.43	$egin{array}{c} 1.21 \ (1.20) \ +0.07 \end{array}$				3.00		

Table 1.—Solar radiation intensities during March 1933—Contd. Madison, Wis.

Sun's zenith distance.													
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon		
mer	75th	Air mass											
	time		Α.	М.				P.M.					
	e.	5.0	4.0	3.0	2.0	1 1.0	2.0	3.0	4.0	5.0	e		
Mar. 1. Mar. 2. Mar. 10. Mar. 15. Mar. 22. Mar. 24. Mar. 27. Means. Departures.	. 86 1. 68 2. 36 2. 49 4. 75		1.04 1.08 1.03 .83	1. 20 1. 10 . 97	1, 50 1, 38 1, 34 1, 08 1, 35	1.61 1.61 1.64	1. 50 1. 24 1. 41 1. 46				mm 2. 36 3. 45 1. 02 1. 60 2. 49 3. 45 3. 81		
•	<u> </u>			Linco	ln, Ne	br.							
Mar. 2	3.00	. 87	1.00	1. 16	1, 16 1, 33	1, 45 1, 38 1, 52 1, 28	1, 22	1.05 1.02	. 90	. 74	3. 30 4. 17 6. 76 3. 63 3. 63		
Mar. 23 Mar. 27	3. 63			1. 10			1.37				3.45		

Departures		+.02	+.04	+.05	01		03			
			3	Blue H	ill, Ma	ass.				
Mar. 5 Mar. 6 Mar. 9	2. 2 1. 5 2. 6				1. 37 1. 30		1, 31 1, 22	0, 89	 	2. 1. 2.
Mar. 10	1. 6 1. 2 2. 0			1. 21 . 85	1, 28 1, 38 , 92		1, 36		 	1. 1. 2.
Mar. 16 Mar. 24 Mar. 25 Mar. 27	2. 2 3. 1 3. 4			1. 15	1. 35 1. 13 1. 17		1. 36 1. 18	1. 17	 	2, 3. 3. 3.
Mar. 30 Means	3. 8 2. 3			1.07	1.30 1.24	1.51	1. 30 1.27		 	2.

¹ Extrapolated.

Table 2.—Average daily totals of solar radiation (direct+diffuse) received on a horizontal surface

	Gram calories per square centimeter												
Week beginning-	Washing- ton	Madison	Lincoln	Chicago	New York	Fresno	Pitts- burgh	Fair- banks	Twin Falls	La Jolla	Gaines- ville	Miami	New Orleans
1933 Feb. 26	cal. 335 582 292 150 378	cal. 254 272 271 275 248	cal. 348 335 370 355 410	cal. 229 249 217 159 253	cal. 248 319 268 211 297	cal. 454 381 402 537 500	cal. 184 237 210 107 265	cal. 172 158 207 205 241	cal. 343 352 348 401 403	cal. 243 299 362 365 361	cal. 381 443 444 385 500	cal. 410 392 475 443 480	cal. 272 288 237 372 328
	Departures from weekly normals												
Feb. 26 Mar. 5 Mar. 12 Mar. 19 Mar. 26	+45 +252 -35 -198 +30	-27 -28 -46 -42 -100	+9 -18 -5 -39 +2	+29 +45 +9 -59 +9	+12 +54 -5 -58 +17	+70 ±0 +2 +79 +2	-2 +33 -7 -119 +14		+43 +39 +19 +53 +17	-90 -39 +13 -9 -42	-30 +45 +45 ±0 -4	+26 -15 +39 -15 +4	
	Accumulated departures on Apr. 2, 1933												
	+2, 394	2, 191	-924	+2, 562	+2, 282	+2, 191	+553		+1,099	-1,568	-5, 131	-567	

Table 3.—Solar-radiation measurements, and determinations atmospheric turbidity factor, \(\beta \), Washington, D.C., March 1933

		_	-			- '			
Date and solar hour angle	Solar alti- tude,	Air mass, m	I cal.	I, cal.	I, cal.	β	Blue- ness of sky	Atmospheric dust particles per cubic centimeter	Notes: Sky light polari zation, P, clouds, etc
1933 Mar. 6 3:54 a	20-01 21-03 21-45 22-28 31-16 31-20	2. 90 2. 77 2. 69 2. 61 1. 92 1. 92	1. 225 1. 244 1. 269 1. 280 1. 285 1. 257	0. 877 .988 .934 .936 .880	0.788 .744 .756 .759 .735	0. 040 . 040 . 038 . 035 . 060 . 070	5	914	P=51.2% Cirri.
Mar. 9 4;18 a	16-34 17-34 24-10 25-10 32-54 33-29 42-08 42-47	3. 48 3. 28 2. 43 2. 34 1. 84 1. 81 1. 49 1. 47	1. 079 1. 105 1. 217 1. 250 1. 340 1. 360 1. 444 1. 463	.791 .797 .881 .879 .918 .921 .921	.665 .667 .707 .712 .746 .747 .735	. 042 . 040 . 045 . 040 . 045 . 045 . 040 . 035	5	27 5	P=48. 2 %
Mar. 27 4:51 a	15-00 15-36 19-40 20-35 26-34 27-18	3. 82 3. 68 2. 95 2. 82 2. 23 2. 17	. 645 . 644 . 753 . 787 . 816 . 832	. 488 . 491 . 569 . 571 . 635 . 638	. 447 . 450 . 516 . 521 . 537 . 540	. 130 . 135 . 165 . 165 . 180 . 180	4	899	Cirri. P=42.6%
Mar. 29 4:57 a	14-24 14-58 19-02 19-56 21-36 22-12 26-51	3. 98 3. 84 3. 05 2. 92 2. 70 2. 63 2. 21	. 909 . 922 1. 029 1. 058 1. 075 1. 106 1, 211	. 688 . 691 . 765 . 769 . 794 . 797	. 591 . 596 . 647 . 650 . 671 . 672 . 700	.065 .065 .075 .065 .078 .072		317	
3:47 a	27-35 43-22 43-56 52-20 52-40 48-19 47-36 38-42 37-40	2. 15 1. 46 1. 44 1. 27 1. 26 1. 34 1. 36 1. 60 1. 63 1. 93 1. 95	1. 236 1. 353 1. 338 1. 424 1. 429 1. 350 1. 306 1. 295 1. 301 1. 218	.856 .888 .891 .935 .938 .950 .963 .884 .887 .860	700 700 700 700 759 759 774 771 694 694 697	.052 .105 .115 .068 .068 .110 .140 .140 .140 .135	6		P = 53.7 %

POSITIONS AND AREAS OF SUN SPOTS

[Communicated by Capt. J. F. Hellweg, Superintendent United States Naval Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of latitude are measured from central meridian, positive west. The north latitude are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column]

	Eastern standard civil time		н	eliograp	hie	Area		Total
Date			Diff. long.	Longi- tude	Lati- tude	Spot	Group	for each day
1933	h.							
Mar. 1 (Naval Observatory)	11	m. 56	-64.0	305. 4	+13.0		123	123
Mar. 2 (Mount Wilson)	12	0	-51.0 -23.0	333. 2	+14.0 +9.0	11	223	234
Mar. 3 (Naval Observatory)	11	34	-37.0	306.3	+13.0		93	93
Mar. 4 (Naval Observatory)	11	35	-24.0	306. 1	+13.0		93	93
Mar. 5 (Perkins Observatory)	16	0	-13.5	301.0	+13.0		90	
			-4.5	310.0	+16.5		90	180
Mar. 6 (Naval Observatory)	11	9	+3.0	307. 0	+13.0		93	93
Mar. 7 (Mount Wilson)	12	35	+16.0	306.0	+14.0		118	118

POSITIONS AND AREAS OF SUN SPOTS-Continued

		Eastern		eliograp	hic	A	rea	Tota area
Date	stand civil t		Diff. long.	Longi- tude	Lati- tude	Spot	Group	for each day
	h.	<i>m</i> .	0	0	•			
Mar. 8 (Naval Observatory)	11	51	$+24.0 \\ +33.0$	301. 2	$+11.0 \\ +15.0$	77	9	8
Mar. 9 (Naval Observatory)	10	5	+37. 0 +46. 0	302. 0 311. 0	+11.0 $+15.0$		9	8
Mar. 10 (Naval Observatory)	11	8	+60.0	311.3	+15.0	77		7
Mar. 11 (Naval Observatory) Mar. 12 (Perkins Observatory)	11 14	49 0		311.7 No spot		77		7
Mar. 13 (Mount Wilson)	îi	3Ŏ		No spot				
Mar. 14 (Mount Wilson)	12	25		No spot				
Mar. 15 (Naval Observatory)	10	58		No spot				
Mar. 16 (Naval Observatory)	11	35		No spot				
Mar. 17 (Naval Observatory)	11	44		No spot				
Mar. 18 (Mount Wilson)	11	0		No spot				
Mar. 19 (Mount Wilson)	10	50		No spot			-	-
Mar. 20 (Mount Wilson)	10	45		No spot				
Mar. 21 (Mount Wilson)	11	0		No spot				Ì
Mar. 22 (Naval Observatory)	13	28 45	-73.0 -58.0	18.8			401 408	40 40
Mar. 23 (Mount Wilson) Mar. 24 (Naval Observatory)	11	40	-35. U -45. 0	20.3	$+5.0 \\ +4.0$		216	21
Mar. 25 (Mount Wilson)	13 11	30	-32.0	21.3	+5.0		234	121
VIAF. 25 (MOUNT WISON)	11	30	+20.0	73.3	+10.0		31	26
Mar. 26 (Naval Observatory)	12	44	-18.0	21. 5	+4.0		247	
viai. 20 (ivavai Observatory)	12	**	+35.0	74.5	+10.0		77	32
Mar. 27 (Mount Wilson)	11	15	-5.0	22. 1	+5.0		200	
Tai. 21 (Modelle Wilson)			+49.0	76. 1	+10.0		44	24
Mar. 28 (Naval Observatory)	11	20	+8.0	21.9	+5.0		185 93	27
f 00 (Novel Observator-)	10	54	$+62.0 \\ +22.0$	75. 9 22. 9	$+10.0 \\ +5.0$		170	17
Mar. 29 (Naval Observatory) Mar. 30 (Naval Observatory)	11	13	+36.0	23. 5	+5.0		123	12
Mar. 31 (Mount Wilson)	11	10	+49.0	23. 3	+4.0		131	13
*1.01.01 (141.00H) ** H30H)		-0			, 1.0		101	•
Mean daily area for March.				i		ŀ	l	12

[Communicated by Capt. J. F. Hellweg, Superintendent United States Naval Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of longitude are measured from central meridian, positive west. The north latitudes are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column]

PROVISIONAL SUN-SPOT RELATIVE NUMBERS FOR MARCH 1933

[Dependent alone on observations at Zurich and its station at Arosa] [Data furnished through the courtesy of Prof. W. Brunner, University of Zurich, Switzerland]

March 1933	Relative numbers	March 1933	Relative numbers	March 1933	Relative numbers
1	10 10 10 13 11	11	7 7 0 0 0	21	0 d12 14 14 Mc24
6 7 8 9 10	a11 11 9 8	16 17 18 19 20	0 0 0 0 0	26	29 26 a28 22 d12

Mean: 30 days = 10.0.

a= Passage of an average-sized group through the central meridian. b= Passage of a large group or spot through the central meridian. c= New formation of a center of activity: E, on the eastern part of the sun's disk; W, on the western part; M, in the central zone. d= Entrance of a large or average-sized center of activity on the east limb.

AEROLOGICAL OBSERVATIONS

[Aerological Division, W. R. Gregg, in charge]

By L. T. SAMUELS

Free-air temperatures during March were above normal at all stations except Chicago and Cleveland. (See table 1.) The departures were only of moderate magnitude. Free-air relative humidities averaged above nor-

mal with departures of small to moderate magnitude in most cases.

There were no important deviations in the monthly resultant winds as compared to the normals.